

What is claimed is:

Subt
A
B
1. A method comprising a multistep process for recovering one or more products from a solution containing one or more components selected from the group consisting of betaine, erythritol, inositol, sucrose, mannitol, glycerol, amino acids and mixtures thereof by using chromatographic separation comprising at least one step, where a weakly acid cation exchange resin is used for the chromatographic separation.

Subt
A
B
2. The method of claim 1 wherein the solution to be treated is a sugar beet derived process solution.

3. The method of claim 2 wherein the sugar beet derived process solution is vinasse, molasses or betaine molasses.

4. The method of claim 1 wherein the product to be recovered is selected from the group consisting of betaine, erythritol, inositol, sucrose, mannitol, glycerol, amino acids and mixtures thereof.

5. The method of claim 1 wherein the product is betaine.

6. The method of claim 1 wherein the product is inositol.

7. The method of claim 1 wherein the product is mannitol.

8. The method of claim 1 wherein at least one column or a part of a column contains a weakly acid cation exchange resin.

9. The method of claim 1 wherein at least one column or a part of a column contains a strongly acid cation exchange resin

10. The method of claim 1 wherein the weakly acid cation exchange resin is an acrylic resin.

11. The method of claim 10 wherein the acrylic resin is derived from the group consisting of methyl acrylate, ethyl acrylate, buthyl acrylate methyl methacrylate and acrylonitrile or acrylic acids or mixtures thereof.

12. The method of claim 11 wherein the resin is in the form selected from the group consisting of Na^+ , K^+ , H^+ , Mg^{2+} and Ca^{2+} .

13. The method of claim 12 wherein the resin is in Na^+ and/or K^+ form.

14. The method of claim 10 wherein the resin is crosslinked with divinyl benzene (DVB).

15. The method of claim 14 wherein the crosslinking degree of the resin is 3 to 8 % by weight.

Sub A3 5

10

15

20

25

30

28. The method of claim 1 wherein the particle size of the weakly acid cation exchange resin is 10 to 2000 μm .

28. The method of claim 1 wherein the particle size of the weakly acidic exchange resin is 10 to 2000 μm .

Sub A

30. The method of claim 1 wherein the pH of the feed solution is from 6 to 11.

5

11

10

10

2

—

1

20

15

25

1

4

30

add B'